## **PCT**

## WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT) (51) International Patent Classification 7: (11) International Publication Number: WO 00/21521 **A2** A61K 31/135, 47/18 (43) International Publication Date: 20 April 2000 (20.04.00) (21) International Application Number: PCT/IB99/01571 (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, EE, ES, FI, 22 September 1999 (22.09.99) (22) International Filing Date: GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, (30) Priority Data: 60/104,024 13 October 1998 (13.10.98) US YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, (71) Applicant (for all designated States except US): PFIZER DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, PRODUCTS INC. [US/US]; Eastern Point Road, Groton, CT 06340 (US). ML, MR, NE, SN, TD, TG). (72) Inventors; and (75) Inventors/Applicants (for US only): HARPER, Nancy, Jane Published [US/US]; 164 Shore Avenue, Groton, CT 06340 (US). Without international search report and to be republished RANADE, Gautam, Ramchandra [IN/US]; 18 Wayne Drive, upon receipt of that report. East Lyme, CT 06333 (US). WELCH, Willard, McKowan

(54) Title: SERTRALINE ORAL CONCENTRATE

[US/US]; 116 Pequot Avenue, Mystic, CT 06355 (US).

(74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., Patent Dept., 235 East 42nd Street, New York, NY 10017 (US).

#### (57) Abstract

The present invention provides an essentially nonaqueous, iliquid pharmaceutical concentrate composition for oral administration containing sertraline or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients. The present invention also provides a use of this concentrate composition to prepare an aqueous solution of sertraline. In addition, the present invention provides a method of using this concentrate composition to treat or prevent a variety of diseases or conditions. Finally, the present invention provides the compound, (1S-cis)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-N-methyl-1-naphthalenamine methanesulfonate.

## SOURCEONE<sup>SM</sup>

For questions regarding this delivery, please contact Customer Services at 800-3-DIALOG.

Please have your DIALOG user number and order number available.

# THE DIALOG CORPORATION THE 11000 Regency Parkway, Suite 10, Cary NC. 27511 - 919-462-8600

This is not an invoice. Charges listed are for informational purposes only and do not include applicable tax, copyright or outside purchase fees or telecommunication charges. Copyrighted material reproduced with permission of copyright owner and further reproduction is prohibited or otherwise subject to restriction.

Document ID: WO200021521A2

For:

PPD, INC.

ATTN: BETSY CRITTENDEN

LEGAL

3151 SOUTH 17TH STREET

WILMINGTON NC 28412

USA

Subject:

DIALOG SOURCEONE

Order Date:

2000/06/26

Number:

200062

User Number:

153275

Subaccount:

DIALOG SOURCEONE

Customer notes:

## FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

ł								
AL	Albania.	25	Spain	LS	Laratho	81	Slovenia .	
AM	Armenia	FI	Finland	LŦ	Lithumia	SK	Siovakia	
AT	Ametria	FR	France	LU	Lexembourg	SN	Senegal	
AU	Australia	GA	Gabon	LY	Lacvia	SZ.	Swaziland	
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Ched	
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo	
BB	Barbedos	GH	Ghuna	MG	Medaguecar	TJ	Tujikistan	
BE	Belgium	GN	Guinca	MK	The former Yugoslav	TM	Terknonistan	
BT	Burkina Paso	GR	Grecce	•	Republic of Macedonia	TR	Turkey	
<b>B</b> G	Bulgaria	Hill	Hungary	ML	Mall	73	Trinkind and Tubero	
53	Sain	==	beland	MEN	Mongolia	CA	Varian	
52	Cracil.	豇	Escael	MER	Materitanic	UG	Ggrade.	
BT	Beiteres	13	iceianci	MW	Malewi	us	United States of Asserting	
CA.	Canada	FF	(Taily	1	Mexico	CT2	Uzbekien	
Œ	Central African Republic		Expen	NE	niger .	VN	Vict Name	
CC	Congo	KE-	Kenya	NL	Netherlands	Yυ	Yagashvin	
CH	Switzerland	KG	Kyrgyzstan	740	riceway	ZW	Zimbelson	
a	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand			
CM	Cameroon		Republic of Korea	PL	Poland			
CN.	China	22	Republic of Ruses	Fi	rostagai			
(ii	Cirbs	N.E.	Katakatan	KŪ	Homania			
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation			
DE	Germany	1.1	Liechtenstein	SD	Syden			
DK	Denmark	LK	Sri Lanks	SE	Sweden			
EE	Estonia	LR	Liberia	SG	Singapore			
E.C.	75 (e. 1)	-	Coord	36	2 members			

5

10

15

20

25

#### SERTRALINE ORAL CONCENTRATE

## FIELD OF THE INVENTION

The present invention provides an essentially nonaqueous, liquid pharmaceutical concentrate composition for oral administration containing sertraline or a pharmaceutically acceptable salt thereof and one or more pharmaceutically acceptable excipients. The present invention also provides a use of this concentrate composition to prepare an aqueous solution of sertraline. In addition, the present invention provides a method of using this concentrate composition to treat or prevent a variety of diseases or conditions. Finally, the present invention provides the compound, (1S-cis)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-N-methyl-1-naphthalenamine methanesulfonate.

#### BACKGROUND OF THE INVENTION

Sertraline is a known compound having the following structure:

Sertraline has the following chemical name: (1S-cls)-4-(3,4-dichlorophenyl)-1,2,3,4-tetrahydro-N-methyl-1-naphthalenamine. Sertraline and its pharmaceutically acceptable acid addition salts, such as the hydrochloride salt, are disclosed in U.S. Patent No. 4,536,518, which issued on 20 August 1985, (hereafter referred to as the '518 patent), which is herein incorporated by reference in its entirety. Another pharmaceutically acceptable salt of sertraline is the mesylate salt.

The '518 patent states that sertraline and derivatives thereof are useful as antidepressant agents. U.S. Patent No. 5,130,338, which issued on 14 July 1992,

refers to the use of sertraline to treat chemical dependencies, including dependencies on alcohol, tobacco and cocaine. U.S. Patent No. 4,962,128, which issued on 9 October 1990, refers to the use of sertraline to treat anxiety related disorders such as panic disorder, obsessive-compulsive disorder, generalized anxiety disorder, phobias, post traumatic stress disorder and avoidant personality disorder. U.S. Patent No. 4,940,731, which issued on 10 July 1990, refers to the use of sertraline to treat premature ejaculation. Published PCT patent application, WO 96/22085, which was published on 25 July 1996, refers to the use of sertraline to treat cancer patients. Published European patent application 0768083, which published on 16 April 1997, refers to the use of sertraline to treat post myocardial infarction patients.

5

10

15

20

25

30

U.S. Patent No. 5,597,826, which issued on 28 January 1997, relates to novel compositions containing a serotonin selective reuptake inhibitor (SSRI), such as sertraline, and an agonist or antagonist of the serotonin 1 (5-HT<sub>1</sub>) receptor and to the use of such compositions for treating or preventing a condition selected from mood disorders, including depression, seasonal affective disorders and dysthmia, anxiety disorders, including generalized anxiety disorder and panic disorder, agoraphobia. avoidant personality disorder; social phobia; obsessive compulsive disorder; posttraumatic stress disorder, memory disorders including dementia, amnestic disorders and age-associated memory impairment; disorders of eating behavior, including anorexia nervosa and bulimia nervosa; obesity; cluster headache; migraine; pain; Alzheimer's disease; chronic paroxysmal hemicrania; headache associated with vascular disorders; Parkinson's disease, including dementia in Parkinson's disease, neuroleptic-induced parkinsonism and tardive dyskinesias; endocrine disorders such as hyperprolactinaemia; vasospasm (particularly in the cerebral vasculature); hypertension; disorders in the gastrointestinal tract where changes in motility and secretion are involved; sexual dysfunction, including premature ejaculation; and chemical dependencies. The foregoing published applications and U.S. patents are herein incorporated by reference in their entireties.

As noted above, sertraline hydrochloride is a selective serotonin reuptake inhibitor (SSRI) for oral administration. (Physicians' Desk Reference (PDR), 52nd ed., Supplement A. pages A250-A255 (1998).) Its mechanism, pharmacokinetics, and metabolism confer safety, once-daily dosage and efficacy in the treatment of depression, obsessive-compulsive disorder and panic disorder. It is approved for the

treatment of one or more of these indications in over 60 countries worldwide. The daily doses for sertraline, expressed as the free base, range from 50-200 mg, increasing in 50 mg increments. A titration dose of 25 mg/day during the initial phase of therapy may be warranted in some indications.

Tablet and capsule formulations of sertraline hydrochloride are commercially available in different countries. Capsular-shaped, scored tablets are available in strengths of 50 and 100 mg and are sold under the brand name, ZOLOFT® Tablets, in the U.S. The capsules are available in some countries in strengths of 50 mg and 100 mg.

5

10

15

20

25

30

The '518 patent discloses that sertraline and related compounds can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically-acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, aqueous suspension, injectable solutions, elixirs, syrups, and the like. According to this patent, when aqueous suspensions and/or elixirs are desired for oral administration, the essential active ingredient therein may be combined with various sweetening, or flavoring agents, coloring matter or dyes and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, otheroi, propylene glycel, glycerin and various like combinations thereof.

However, development of an oral liquid dosage form of sertraline has been complicated by the objectionable bitter taste and astringency sensation imparted by the drug in liquid form. Thus, direct ("ready-to-use") oral liquid solutions or suspensions of sertraline, such as those described in the '518 patent above, have an objectionable taste, despite the inclusion of a variety of taste-masking or flavoring agents.

An oral liquid dosage form of sertraline with acceptable taste would be a valuable addition to the existing formulations, providing greater choice for both the prescriber and the patient. This is of importance with regard to the issue of non-compliance with treatment, which is believed to affect up to 50% of outpatients and appears to be a particular problem with elderly, pediatric and psychiatric patients (B. Blackwell, Drug Therapy: Patient Compliance, N.Engl.J.Med. 1973, 289(5):249-52). By virtue of being easier to swallow, an oral liquid dosage form of sertraline would offer an alternative to those patients who dislike or have difficulty swallowing

tablets or capsules, and would therefore be of considerable benefit in treating those who may be non-compliant for those reasons. Therefore, there is a need in the art for an alternative liquid dosage form of sertraline that has acceptable taste and properties.

Oral concentrate drug products are known in the art and are commercially available. However, these concentrates are conventionally aqueous. For example, NAVANE® Concentrate contains the active ingredient, thiothixene hydrochloride, and the inert ingredients of alcohol, cherry flavor, dextrose, passion fruit flavor, sorbitol solution and water. (Physicians' Desk Reference (PDR), 52nd ed., pages 2192-2193 (1998).) SINEQUAN® Concentrate contains the active ingredient, doxepin hydrochloride, and the inert ingredients of glycerin, methylparaben, peppermint oil, propylparaben and water. (Physicians' Desk Reference (PDR), 52nd ed., pages 2203-2204 (1998).) TRILAFON® Concentrate contains the active ingredient, perphenazine, and the inert ingredients of alcohol, citric acid, flavors, menthol, sodium phosphate, sorbitol, sugar and water. (Physicians' Desk Reference (PDR), 52nd ed., pages 2666-2668 (1998)).

## **SUMMARY OF THE INVENTION**

The present invention particularly provides:

5

10

15

20

25

30

A pharmaceutical composition which comprises:

an essentially nonaqueous, liquid concentrate for oral administration comprising an amount of sertraline or a pharmaceutically acceptable salt thereof and one or more essentially nonaqueous pharmaceutically acceptable excipients; wherein at least one of the excipients is liquid. Preferably, the pharmaceutically acceptable salts of sertraline are the hydrochloride salt and the mesylate salt.

More particularly, the present invention provides this composition wherein the concentrate contains sertraline hydrochloride, ethanol and glycerin. More specifically, in this concentrate, the scrtraline hydrochloride is present in an amount of about 15 to about 30 mg/ml and the ethanol and glycerin are present at about 8 to about 20% ethanol (by weight) in glycerin.

The present invention also provides this composition wherein the concentrate contains sertraline hydrochloride, ethanol, glycerin and menthol. More specifically, in

this concentrate, the sertraline hydrochloride is present in an amount of about 15 to about 30 mg/ml, the ethanol and glycerin are present at about 8 to about 20% ethanol (by weight) in glycerin, and the menthol is present in an amount of about 0.01 to about 5.0 mg/ml.

Most particularly, the present invention provides this composition wherein the concentrate contains sertraline hydrochloride, glycerin, ethanol, butylhydroxytoluene (BHT) and menthol. More specifically, each ml of this concentrate contains about 22.4 mg of sertraline hydrochloride, about 151 mg of ethanol, about 0.50 mg of menthol, about 0.10 mg of BHT, and a sufficient quantity of glycerin (which is about 1011 mg) to produce about 1 ml.

5

10

15

20

25

30

The present invention also provides the use of this essentially nonaqueous, liquid concentrate of sertraline or a pharmaceutically acceptable salt thereof to prepare an aqueous solution of sertraline which comprises: diluting the concentrate in an aqueous diluent prior to oral administration. Preferably, the pharmaceutically acceptable salts of sertraline are the hydrochloride salt and the mesylate salt.

The present invention also provides a method of treating or preventing diseases or conditions which are caused by disorders of the serotonergic system which comprises: a) diluting this concentrate in an aqueous diluent; and b) orally administering the resulting aqueous solution to a patient in need thereof. In addition, the present invention provides this method for treating or preventing diseases or conditions, such as the following: depression, anorexia, chemical dependencies, anxiety-related disorders (such as panic disorder, obsessive-compulsive disorder, generalized anxiety disorder, phobias, post traumatic stress disorder and avoidant personality disorder), premature ejaculation, cancer and post myocardial infarction.

The present invention also provides a process for making this essentially nonaqueous, liquid concentrate of sertraline or a pharmaceutically acceptable salt thereof.

Finally, the present invention provides the compound, (1S-cis)-4-(3,4-dichlorophenyi)-1,2,3,4-tetrahydro-N-methyi-1-naphthalenamine methanesuifonate.

## DETAILED DESCRIPTION OF THE INVENTION

"Essentially nonaqueous" when used herein refers to the amount of water that is present in the final drug product, which amount is consistent with the amount of

water potentially contributed by the active ingredient and/or by the excipients; however, no water is directly added to the final drug product. About 10% is the upper limit of the amount of water that may be present in the oral concentrate of the present invention. Typically, the amount of water that is present in the oral concentrate of the present invention ranges from about 0.8% to about 5.2% of the composition. Most typically, the amount of water that is present ranges from about 0.8% to about 2.0% of the composition.

"Concentrate" when used herein refers to a strong solution provided for dilution before use. (Butterworths Medical Dictionary, 2nd edition, Butterworths, London - Boston 1978, pp. 399-400.)

.10

15

20

25

30

"Pharmaceutically acceptable excipient," as used herein, means a pharmacologically inactive ingredient which is used to formulate a drug product and which is readily known and available in the pharmaceutical arts. Examples of such excipients are given below.

"Pharmaceutically acceptable preservative," as used herein, means an additive or excipient that preserves the quality of the formulation from a chemical or microbiological standpoint. Examples of preservatives are given below.

"Serctonergic system" when used herein refers to the neuronal precess whereby serotonin is released by a pre-synaptic cell upon excitation and crosses the synapse to stimulate or inhibit the post-synaptic cell.

The sertraline oral concentrate of the present invention is a new dosage form of sertraline intended as a multidose product for oral use only. The oral concentrate is an essentially non-aqueous, optionally flavored solution of sertraline, preferably as the hydrochloride salt, which is intended to be diluted prior to administration.

The oral concentrate of the present invention is advantageous because, inter alia, it provides convenience in measuring different doses, which are needed for certain indications, and acceptable taste upon administration, as well as good physical/chemical stability characteristics throughout the product's shelf-life and use interval. Also, since the concentrate of the present invention is a solution, it is preferred over a suspension for ease of manufacture and optimal control of dosing homogeneity.

The active ingredient in the concentrate of the present invention is sertraline, which may be present as its free base or as its pharmaceutically acceptable sait.

Such salts include, for example, the hydrochloride salt and the methanesulfonate salt (also known as the mesylate salt). In the concentrate of the present invention, the hydrochloride salt is preferred.

The preparation of sertraline and its hydrochloride salt is described in the '518 patent referenced above, particularly in Example 2 of that patent. The mesylate salt is prepared by reacting the sertraline free base with methanesulfonic acid at room temperature in a solvent, such as ethyl acetate, methanol, ethanol and isopropanol. The mesylate salt has favorable properties, such as solubility in polar vehicles and good chemical stability.

5

10

15

20

25

30

Sertraline hydrochloride may be present in the concentrate of the present invention in an amount of about 1 mg/ml up to about 88 mg/ml of the composition. Preferably, it is present in an amount of about 15 to about 30 mg/ml of the composition. Most preferably, about 20 mg sertraline per ml, which is equivalent to about 22.4 mg/ml as the hydrochloride (assuming a theoretical stoichiometric potency of 89.4%), is present. The quantity of sertraline hydrochloride needed to produce a concentration of about 20 mg sertraline per ml is adjusted based on the actual potency of the drug substance lot to be used.

One or more pharmaceutically acceptable excipients are present in the concentrate of the present invention. Examples of such excipients include the following: ethanol, glycerin (also referred to as glycerol or glycerine), propylene glycol and polyethylene glycols. All of these excipients are well known in the pharmaceutical arts and are commercially available.

From among these exciplents, binary combinations of ethanol and glycerin provide maximum solubilization of the sertraline hydrochloride drug substance and are preferred. In these combinations, the amount of ethanol that is present is about 8% to about 50% (w/w) in glycerin; preferably, about 8 to about 20% (w/w) ethanol in glycerin is present; most preferably, about 10 to about 14% (w/w) ethanol in glycerin is present. The ethanol that may be used in the concentrate of the present invention is any compendial grade of ethanol, including ethanol (95%) and anhydrous ethanol.

The concentrate of the present invention optionally contains one or more flavoring agents or taste-masking flavors. Examples of candy-type flavoring agents include mint, citrus and fruit flavoring agents. The mint flavoring agents, such as peppermint, spearmint and menthol (also referred to as levomenthol), are preferred.

More specifically, from among a variety of mint flavoring agents, menthol is preferred and may be present in an amount of about 0.01 to about 5.0 mg/ml of the composition; preferably, menthol is present in an amount of about 0.5 mg/ml of the composition.

5

10

15

20

25

30

The concentrate of the present invention also optionally contains one or more pharmaceutically acceptable preservatives, which preserve the quality of the concentrate from a chemical or microbiological standpoint. Preservatives, such as antioxidants, metal chelators, metal complexing agents and antimicrobial agents, may be used in the concentrate of the present invention. These preservatives may be used in the concentrate, both alone and in various combinations (e.g., antioxidant(s) in combination with a metal complexing agent), to improve product stability. When the concentrate of the present invention contains ethanol, it may be appropriately microbiologically self-preserved.

Examples of preservatives, which may be present in the concentrate of the present invention, include the following: butylhydroxytoluene (BHT, also referred to as butylated hydroxytoluene), butylated hydroxyanisole (BHA), propyl gallate, ascorbic acid, ascorbyl palmitate, sodium metabisulite, sodium bisulfite, sodium thiosulfate, sodium hydroxide, cystiene, ethylenediamine tetraacetic acid (EDTA) or its salts (such as the sodium salt), citric acid, triethanolamine, thioglycerol, methylparaben and propylparaben. Also, various combinations of these preservatives may be present in the concentrate of the present invention, such as the following: disodium EDTA and BHA; disodium EDTA, BHA and BHT; disodium EDTA and propyl gallate; and BHA and BHT.

The amount of preservative that may be present in the concentrate of the present invention ranges from about 0.01 to about 10.0 mg/ml of the composition. In the concentrate of the present invention, butylhydroxytoluene (BHT) is preferred and may be present in an amount of about 0.01 to about 0.5 mg/ml of the composition; preferably, it is present in an amount of about 0.01 to about 0.2 mg/ml of the composition; more preferably, it is present in amount of about 0.01 to about 0.13 mg/ml of the composition; most preferably, it is present in an amount of about 0.05 to about 0.11 mg/ml of the composition.

The sertraline oral concentrate of the present invention is intended to be diluted in a suitable diluent or beverage prior to administration. For example, doses

of the oral concentrate (equivalent to about 25 mg and about 200 mg of sertraline) are mixed with approximately 4 oz. (about 120 ml or 1/2 cup) of a suitable diluent or beverage. The following beverages may be used for the preparation and administration of the dose: water, orange juice, ginger ale, lemon-lime soda, lemonade, sugared tap water, cranberry juice, grapefruit juice, tomato juice, pineapple juice, prune juice, orange drink and Gatorade® (beverage, not powder). Water, orange juice, ginger ale, lemon-lime soda and lemonade are preferred. The proper dose of sertraline oral concentrate may be most conveniently added to these beverages by means of a dropper calibrated at, for example, 25 and 50 mg (as sertraline).

The manufacture of sertraline oral concentrate utilizes conventional pharmaceutical equipment and processes. For example, the manufacturing process for sertraline hydrochloride oral concentrate involves dissolving the sertraline hydrochloride drug substance in a solution of glycerin, ethanol, and optionally other excipients, such as butylhydroxytoluene (BHT) and menthol, followed by clarifying filtration, if desired, and filling into containers such as bottles, as appropriate.

10

15

20

25

Preferably, to facilitate BHT incorporation, this component is first dissolved in a portion of the ethanoi, which is then added to the main compounding tank containing the glycerin and the remainder of the ethanoi. Menthol and sertraline hydrochloride are then sequentially added and mixed to dissolve. The sertraline hydrochloride is intentionally added to the solution last to optimize the protective effect of the antioxidant (BHT). The compounded solution is passed through a filter, and the final product is filled into appropriate containers, such as bottles. The final product may be supplied in a variety of presentations, such as a 60 ml presentation, which is packaged in, for example, a 60 ml, amber glass bottle with a child-resistant closure.

#### **EXAMPLES**

-10-

EXAMPLE 1 Concentrate solution (20 mg sertraline/ml) with preservative and menthol flavoring:

4		

Ingredients	mg/ml	% w/v (g/100ml)	mg/60ml
Sertraline HCI	22.37	2.237	1342.3
Ethanol (95%)	150.68	15.068	9040.8
Menthol	0.50	0.050	30.0
ВНТ	0.10	0.010	6.0
Glycerin	1010.81	101.081	60648.6
Total	1184.46	118.446	71067.7

BHT, menthol and drug were dissolved in a mixture of glycerin and ethanol. The resulting mixture may be blanketed with nitrogen due to the ethanol content and associated potential flammability. The resulting mixture may be filtered, if desired.

## **EXAMPLE 2**

One ml of a nonaqueous solution containing 1-23 mg sertraline HCl/ml:

Sertraline Hydrochloride: 1 - 23 mg

Polyethylene glycol 300 (PEG-300): sufficient quantity to produce 1 ml

15 Dissolve the drug substance in the PEG with stirring.

## EXAMPLE 3

One ml of a nonaqueous solution containing 1 - 64 mg sertraline HCI/ml:

Sertraline Hydrochloride: 1 - 64 mg

Propylene Glycol: sufficient quantity to produce 1 ml

20 Dissolve the drug substance in the propylene giycoi with stirring.

## EXAMPLE 4

One ml of a nonaqueous solution containing 1 - 17 mg sertraline HCl/ml:

Sertralino Hydrochlorido: 1 17 mg

Ethanol: sufficient quantity to produce 1 ml

25 Dissolve the drug substance in the ethanol with stirring.

## **EXAMPLE 5**

One ml of a nonaqueous solution containing 1 - 78 mg sertraline HCl/ml:

Sertraline Hydrochloride: 1 - 78 mg

Ethanoi: 100 - 500 mg

5 Glycerin: sufficient quantity to produce 1 ml

Dissolve the drug substance in a mixture of ethanol and glycerin with stirring.

**EXAMPLE 6** 

1000 ml concentrate solution (20 mg sertraline/ml) with peppermint flavoring:

Sertraline HCI: 22.4 g

10 Ethanol (95%): 127 g

Natural Peppermint Stick flavoring: 0.01 - 1.7 g

Glycerin: 852 - 872 g or sufficient quantity to produce 1000 ml

Dissolve the natural peppermint stick flavoring and drug in a mixture of ethanol and glycerin.

15 EXAMPLE 7

1000 ml concentrate solution (20 mg sertraline/ml) with menthol flavoring:

Sertraline HCI: 22.4 g

Ethanol (95%): 127 q

Menthol: 0.01 - 1.7 g

20 Glycerin: 852 - 872 g or sufficient quantity to produce 1000 ml

Dissolve the menthol and drug in a mixture of ethanol and glycerin.

**EXAMPLE 8** 

1000 ml concentrate solution (20 mg sertraline/ml) with preservatives and peppermint

flavoring:

25 Sertraline HCl: 22.4 g

Ethanol, anhydrous: 127 g

Natural Peppermint Stick flavoring: 1.7 g

Methylparaben: 0.6 g

Propylparaben: 0.3 g

30 Glycerin: 852 - 872 g or sufficient quantity to produce 1000 mi

The natural peppermint stick flavoring, parabens and drug were dissolved in a

mixture of ethanol/glycerin.

## **EXAMPLE 9**

1000 ml concentrate solution (20 mg sertraline/ml) with preservatives and no

flavoring:

Sertraline HCI: 22.4 g

5 Ethanol, anhydrous: 127 g

Methylparaben: 0.6 g Propylparaben: 0.3 g

Glycerin: 852 - 872 g or sufficient quantity to produce 1000 ml

The parabens and drug were dissolved in a mixture of ethanol/glycerin.

## 10 EXAMPLES 10-30

The following examples describe various 1000 ml concentrate solutions of sertraline hydrochloride (20 mg sertraline/ml) with preservatives and menthol flavoring:

**EXAMPLE 10** 

Sertraline HCI: 22.7 g

15 Ethanol, 95%: 151.0 g

Menthol: 0.5 g Citric acid: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 11** 

20 Sertraline HCl: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Propyi gallate: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

25 EXAMPLE 12

Sertraline HCI: 22.7 g

Ethanol, 95%: 151.0 g

Menthal: 0.5 g

BHA: 0.1 g

30 Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 13** 

Sertraline HCl: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Sodium metabisulfite: 0.5 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

5 EXAMPLE 14

Sertraline HCl: 22.7 g Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Ascorbyl palmitate: 0.1 g

10 Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 15** 

Sertraline HCl: 22.7 g Ethanol, 95%: 151.0 g

Menthol: 0.5 g

15 Sodium bisulfite: 0.5 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

EXAMPLE 16

Sertraline HCl: 22.7 g Ethanol, 95%: 151.0 g

20 Menthol: 0.5 g

Triethanolamine: 0.1 - 0.5 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 17** 

Sertraline HCI: 22.7 g

25 Ethanol, 95%: 151.0 g

Menthol: 0.5 g

BHT: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

EXAMPLE 18

30 Sertraline HCl: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

PCT/IB99/01571

1N sodium hydroxide: 0.03 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 19** 

Sertraline HCl: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Disodium EDTA: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 20** 

Sertraline HCl: 22.7 g 10

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Ascorbic acid: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 21** 15

Sertraline HCI: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Thioglyceroi: 1.0 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml 20

EXAMPLE 22

Sertraline HCI: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Sodium thiosulfate: 0.5 g 25

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 23** 

Sertraline HCl: 22.7 g

Ethanoi, 95%: 151.0 g

30 Menthol: 0.5 g

BHT: 0.1 g

BHA: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

-15-

**EXAMPLE 24** 

Sertraline HCl: 22.7 g Ethanol, 95%: 151.0 g

5 Menthol: 0.5 g

BHT: 0.1 g BHA: 0.1 g

Citric acid: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

10 EXAMPLE 25

Sertraline HCl: 22.7 g Ethanol, 95%: 151.0 g

Menthol: 0.5 g

BHT: 0.1 g

15 BHA: 0.1 g

Propyl gallate: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 26** 

Sertraline HCl: 22.7 g

20 Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Disodium EDTA: 0.1 g

Sodium metabisulfite: 0.5 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

25 EXAMPLE 27

Sertraline HCl: 22.7 g Ethanol, 95%: 151.0 g

Menthol: 0.5 g Cystiene: 1.0 g

30 Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

EXAMPLE 26

Sertreiine HCi: 22.7 g

WO 00/21521 -16- PCT/IB99/01571

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

BHT: 0.1 g

BHA: 0.1 g

5 Triethanolamine: 0.1 - 0.5 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

**EXAMPLE 29** 

Sertraline HCi: 22.7 g

Ethanol, 95%: 151.0 g

10 Menthol: 0.5 g

Disodium EDTA: 0.1 g

BHT: 0.1 g

BHA: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

15 EXAMPLE 30

Sertraline HCI: 22.7 g

Ethanol, 95%: 151.0 g

Menthol: 0.5 g

Disodium EDTA: 0.1 g

20 BHA: 0.1 g

Glycerin: 1010.7 g or sufficient quantity to produce 1000 ml

For Examples 10-30 above, the menthol and preservative were dissolved in ethanol.

Glycerin was added and dispersed. The drug was added and dissolved.

## CLAIMS

1. A pharmaceutical composition which comprises:

15

an essentially nonaqueous, liquid concentrate for oral administration comprising an amount of sertraline or a pharmaceutically acceptable salt thereof and one or more essentially nonaqueous pharmaceutically acceptable excipients; wherein at least one of the excipients is liquid.

- 2. The composition of claim 1 wherein the pharmaceutically acceptable salt of sertraline is the hydrochloride salt or the mesylate salt.
- 3. The composition of claim 1 wherein the excipients are selected from the group consisting of ethanol, glycerin, polyethylene glycol and propylene glycols.
  - 4. The composition of claim 3 wherein the excipients are ethanol and glycerin.
  - 5. The composition of claim 1 wherein the concentrate comprises sertraline hydrochloride in an amount of about 15 to about 30 mg/ml and wherein the excipients are ethanol and glycerin in an amount of about 8 to about 20% ethanol (by weight) in glycerin.
  - 6. The composition of claim 5 wherein the concentrate further comprises one or more flavoring agents and one or more pharmaceutically acceptable preservatives.
- The composition of claim 6 wherein the flavoring agents are selected from the group consisting of peppermint, spearmint and menthol; and wherein the
   preservatives are selected from the group consisting of butylhydroxytoluene, butylated hydroxyanisole, propyl gallate, ascorbic acid, ascorbyl palmitate, sodium metabisulite, sodium bisulfite, sodium thiosulfate, sodium hydroxide, cystiene, ethylenediamine tetraacetic acid or salts thereof, citric acid, triethanolamine, thioglycerol, methylparaben and propylparaben.
- 25 8. The composition of claim 7 wherein the flavoring agent is menthol and wherein the preservative is butylhydroxytoluene.
  - 9. The composition of claim 8 wherein each ml of the concentrate comprises about 22.4 mg of sertraline hydrochloride, about 151 mg of ethanol, about 0.50 mg of menthol, about 0.10 mg of butylhydroxytoluene, and about 1011 mg of glycerin.
- 30 10. The compound, (1S-cis)-4-(3,4-dichlorophenyi)-1,2,3,4-tetrahydro-N-methyi-1-naphthaienamine methanesulfonate.
  - 11. Use of an essentially nonaqueous, liquid concentrate of sertraline or a pharmaceutically acceptable salt thereof of claim 1 to prepare an aqueous solution of

sertraline which comprises diluting the concentrate in an aqueous diluent prior to oral administration.

- 12. The use of claim 11 wherein the pharmaceutically acceptable salt of sertraline is the hydrochloride salt or the mesylate salt.
- 5 13. The use of claim 11 wherein the diluent is selected from the group consisting of water, orange juice, ginger ale, lemon-time soda and lemonade.